

Passage of Ingested *Mansonella ozzardi* (Spirurida: Onchocercidae) Microfilariae Through the Midgut of *Aedes aegypti* (Diptera: Culicidae)

JEFFERSON A. VAUGHAN,^{1,2} JEFFREY A. BELL,¹ MICHAEL J. TURELL,³ AND DAVE D. CHADEE⁴

J. Med. Entomol. 44(1): 111–116 (2007)

ABSTRACT When virus and microfilariae are ingested concurrently by a mosquito, microfilariae (mf) may penetrate the mosquito midgut and introduce virus directly into the mosquito hemocoel, allowing mosquitoes to become infectious much sooner than normal and enhancing transmission of viruses by mosquitoes. *Mansonella ozzardi* (Manson) is a benign filarial nematode parasite of humans in Latin America and is transmitted by black flies (Diptera: Simuliidae) and biting midges (Diptera: Ceratopogonidae). Because *M. ozzardi* and dengue are sympatric, we wanted to know whether *M. ozzardi* mf had the ability to penetrate the midgut of *Aedes aegypti* (L.) (Diptera: Culicidae) and thus play a potential role in the enhancement of dengue transmission. To test this, the F1 progeny from locally collected *Ae. aegypti* were fed on *M. ozzardi*-infected human males in an endemic village in northern Trinidad. Mosquitoes were dissected at various times after feeding and examined for mf in the midguts and thoraces. Microfilariae penetrated the midguts of 43% of 63 mosquitoes that ingested mf. Overall, 11% of mf penetrated the midgut by 17 h after being ingested. The intensity of midgut penetration was positively correlated to the numbers of mf ingested. Because midgut penetration is a key requirement for mf enhancement to occur, the potential exists that *M. ozzardi* could be involved in the enhancement of dengue virus transmission.

KEY WORDS *Mansonella ozzardi*, *Aedes aegypti*, dengue, microfilaria, transmission

Concurrent ingestion of microfilariae (mf) and arboviruses by vector arthropods has been shown to result in significantly more efficient transmission of the virus than when the same dose of virus was ingested alone (Mellor and Boorman 1980; Turell et al. 1984, 1987; Zytoon et al. 1993; Vaughan and Turell 1996; Vaughan et al. 1999). This has been referred to as microfilarial enhancement of arboviral transmission. As part of their normal developmental cycle in a vector, most mf species penetrate the vector midgut after being ingested and migrate to a preferred site of development (e.g., flight muscle and fat body). If a microfilaremic host is also viremic when fed upon by a vector, then there is the possibility that penetration of the midgut by mf will allow some of the ingested virus to enter directly into the vector hemocoel. The facilitated movement of virus directly into the hemocoel can have two important epidemiological consequences.

First, it can increase vector competence by allowing virus to bypass both midgut infection and midgut escape barriers (Chamberlain and Sudia 1961, Kramer et al. 1981), transforming otherwise incompetent vector species into competent vector species and thus increasing the number of vector species involved in an arbovirus transmission cycle. Second, mf enhancement can accelerate arboviral development within a vector and shorten the time required for a virus-exposed mosquito to become infectious (i.e., shorten the extrinsic incubation period [EIP]). Because the EIP affects transmission exponentially (MacDonald 1952), small reductions in EIP can lead to large increases in vectorial capacity, even within natural arbovirus-vector systems.

Most laboratory studies of mf enhancement have used various arboviruses with *Brugia* spp. mf and *Aedes/Ochlerotatus* spp. mosquitoes as experimental model systems (Turell et al. 1984, 1987; Vaughan and Turell 1996; Vaughan et al. 1999). But the concept of mf enhancement could apply, in theory, to almost any filarial-vector system (Mellor and Boorman 1980) where filarial and arboviral infections are sympatric. An important consideration is whether the mf can penetrate the midgut and enter the hemocoel and thus allow the virus to bypass the midgut. Further development of the parasite is irrelevant, i.e., a mosquito species does not have to be a competent vector of

The opinions, interpretations, conclusions, and recommendations are those of the authors and are not necessarily endorsed by the U.S. Army.

¹ Department of Biology, University of North Dakota, Grand Forks, ND 58202-9019.

² Corresponding author, e-mail: jefferson_vaughan@und.nodak.edu.

³ Virology Division, U.S. Army Medical Research Institute of Infectious Diseases, Fort Detrick, MD 21702.

⁴ Department of Life Sciences, University of the West Indies, St. Augustine, Trinidad and Tobago, West Indies.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1 JAN 2007		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Passage of ingested Mansonella ozzardi (Spirurida:Onchocercidae) microfilariae through the midgut of Aedes aegypti (Diptera:Culicidae). Journal of Medical Entomology 44:111-116				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Vaughan, JA Bell, JA Turell, MJ Chadee, DD				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) United States Army Medical Research Institute of Infectious Diseases, Fort Detrick, MD				8. PERFORMING ORGANIZATION REPORT NUMBER TR-06-072	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT When virus and microfilariae are ingested concurrently by a mosquito, microfilariae (mf) may penetrate the mosquito midgut and introduce virus directly into the mosquito hemocoel, allowing mosquitoes to become infectious much sooner than normal and enhancing transmission of viruses by mosquitoes. Mansonella ozzardi (Manson) is a benign filarial nematode parasite of humans in Latin America and is transmitted by black flies (Diptera: Simuliidae) and biting midges (Diptera: Ceratopogonidae). Because M. ozzardi and dengue are sympatric, we wanted to know whether M. ozzardi mf had the ability to penetrate the midgut of Aedes aegypti (L.) (Diptera: Culicidae) and thus play a potential role in the enhancement of dengue transmission. To test this, the F1 progeny from locally collected Ae. aegypti were fed on M. ozzardi-infected human males in an endemic village in northern Trinidad. Mosquitoes were dissected at various times after feeding and examined for mf in the midguts and thoraces. Microfilariae penetrated the midguts of 43% of 63 mosquitoes that ingested mf. Overall, 11% of mf penetrated the midgut by 17 h after being ingested. The intensity of midgut penetration was positively correlated to the numbers of mf ingested. Because midgut penetration is a key requirement for mf enhancement to occur, the potential exists that M. ozzardi could be involved in the enhancement of dengue virus transmission.					
15. SUBJECT TERMS Mansonella ozzardi, Aedes aegyptii, dengue, microfilararae, enhancement, transmission					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

filariosis for mf enhancement to occur. This was illustrated in a study examining the ability of co-ingested mf of *Brugia malayi* to affect the ability of *Ochlerotatus taeniorhynchus* (Wiedemann) (Diptera: Culicidae) transmit Venezuelan equine encephalomyelitis virus (family *Togaviridae*, genus *Alphavirus*, VEEV) (Vaughan et al. 1999). A 30-fold increase in VEEV transmission rate was observed in the mosquitoes ingesting both agents compared with mosquitoes ingesting VEEV alone, yet ingested *B. malayi* mf never developed to larval (L3) stage within *Oc. taeniorhynchus* and were essentially all dead by 48 h after ingestion (J.A.V., unpublished data). Potentially, there could be innumerable combinations of parasite-vector-virus in nature where mf enhancement could apply, including combinations with mf not normally associated with mosquitoes.

Dengue is the most important mosquito-borne viral disease of humans and involves cyclical transmission of the virus between humans and mosquito vectors [primarily *Aedes aegypti* (L.)]. There are also several human filarial diseases that are sympatric with dengue, including lymphatic filariasis (*Wuchereria bancrofti*, *Brugia malayi*, and *Brugia timori*), loiasis (*Loa loa*), onchocerciasis (*Onchocerca volvulus*), and mansonielliasis (*Mansonella ozzardi*, *Mansonella perstans*, and *Mansonella streptocerca*). To understand the potential that these human filarial infections may have on the mf enhancement of dengue transmission, it is important to understand the ability of these mf species to penetrate the midgut of *Ae. aegypti*. It is well established that *Wuchereria* and *Brugia* mf can readily penetrate the midgut of *Ae. aegypti* mosquitoes (Ramachandran and Zaini 1967, McGreevy et al. 1982, Lowichik and Lowrie 1988, Zielke 1992, Calheiros et al. 1998). Indeed, laboratory studies have shown that co-ingestion of *B. malayi* mf and dengue 2 virus can enhance virus transmission by *Ae. aegypti* (Turell et al. 1987). Less is known about the ability of the other, nonmosquito-borne human filarial species to penetrate the midgut of *Ae. aegypti* mosquitoes. *Mansonella ozzardi* is a benign filarial infection of humans throughout tropical Latin America that is transmitted by black flies (Diptera: Simuliidae: *Simulium* spp.) and/or biting midges (Diptera: Ceratopogonidae: *Culicoides* spp.) (Nelson and Davies 1976, Nathan 1978, Shelley et al. 1980, Shelley and Coscaron 2001). Although not generally associated with mosquitoes, *M. ozzardi* occurs in areas where several mosquito-borne arboviruses are transmitted, including dengue virus. In this report, we describe the ability of *M. ozzardi* mf to penetrate the midgut of a co-indigenous strain of *Ae. aegypti* mosquitoes.

Materials and Methods

Study Site. The research was conducted on-site within the village of Blanchisseuse (10° 78' N, 61° 31' W), a discrete rural community on the coastal foothills of Trinidad's forested Northern Range Mountains. The village is endemic for *M. ozzardi* and has been a study site for *Mansonella* research for many years

(Nathan et al. 1982, Chadee et al. 1995, Gonzalez et al. 1999).

Mosquitoes. Engorged *Ae. aegypti* females were collected on-site in Blanchisseuse and returned to the Insect Control Division, Ministry of Public Health, in Saint Joseph, Trinidad and Tobago, where the females were allowed to lay eggs and larvae were reared to adults. To safeguard against the potential of using mosquitoes that may have been vertically infected with dengue or other flavivirus, the parents of the F1 mosquitoes ($n = 115$) were frozen and shipped overnight to the University of North Dakota where they were tested by polymerase chain reaction (PCR) against flavivirus RNA by using the primer sets and methodology described by Scaramozzino et al. (2001). The RNA extracted from locally caught *Culex tarsalis* (Coquillett) (Diptera: Culicidae) infected with West Nile virus (Bell et al. 2005) served as positive controls. All *Ae. aegypti* parent mosquitoes tested negative, ensuring that all F1 mosquitoes used in this trial were derived from flavivirus-free stock.

Mosquito Blood Feeding. Before this research was conducted, the study protocol received approval from the Institutional Ethics Committee of the Ministry of Health, Trinidad and Tobago and the Human Subjects Research Review Board of the University of North Dakota. Potential subjects (males older than 19-yr old) were recruited from villagers known from previous surveys to harbor *M. ozzardi* infections. Two days before mosquito blood feeding, 200 μ l of blood was obtained from each of six subjects via fingerprick, spread on a microscope slide, air-dried overnight, and stained with Giemsa stain, and the total number of mf on each slide was counted to determine absolute density of mf in the peripheral blood for each person. Mosquitoes were allowed 15–20 min to feed from a screened cage, based on procedures approved by the two Institutional Review Boards. Most of the mosquito feedings were conducted in the evening (2030–2230 hours). Two of volunteers with the highest microfilaremias allowed mosquitoes to feed on their arms the following afternoon (1500–1700 hours) to test for possible mf periodicity that might affect the level of mf uptake by a diurnal feeding mosquito such as *Ae. aegypti*.

Assessment of Microfilarial Ingestion and Penetration. Engorged mosquitoes were held at ambient temperature. Beginning 1 h after the feeding, three people (two dissecting mosquitoes; one counting mf) collected paired data for each mosquito (i.e., mf counts for bloodmeal and carcass). Mosquitoes were aspirated into ethanol, transferred to saline, and individual midguts were carefully excised with fine-tipped jeweler's forceps, taking care not to rupture the midgut during the process. Only "clean" dissections where midguts were excised intact were included in the data set. Individual midguts were transferred to a glass slide, compressed with a coverslip, and examined for mf at 100–400 \times magnification. The carcasses were thoroughly teased apart with needles and similarly examined for mf. Microfilariae were relatively easy to find when they were alive and wriggling, but dead or

Table 1. Ingestion of *M. ozzardi* mf by endemic F1 *Ae. aegypti* mosquitoes fed on the forearms of six different microfilaremic adult male human volunteers, Blanchisseuse Trinidad, 2004

Volunteer no. and age (yr)	Microfilaremia (no. mf/200 μ l)	Expected no. mf ingested/mosquito ^a	Actual no. mf ingested/mosquito ^b	% ingesting mf (no. examined)
2 (90)	394	5.9–9.8	8.6 (5.9, 12.4)	94 (35)
3 (62)	85	1.3–2.1	1.9 (1.1, 2.9)	79 (24)
4 (61)	73	1.1–1.8	1.6 (0.4, 3.6)	75 (8)
5 (58)	12	0.2–0.3	0	0 (4)
6 (88)	10	0.1–0.2	0.3 (0, 1.1)	40 (5)
1 (62)	3	0.04–0.07	0.3 (0, 0.7)	37 (8)

^a Based on the mf density of volunteer blood per microliter times the average range of bloodmeal volumes measured for *Ae. aegypti* (=3–5 μ l) as reported in Clements 1992.

^b Geometric mean (upper, lower 95% CL).

moribund mf often required careful inspection. For each mosquito, mf were counted as either being contained within the bloodmeal or in the hemocoel. The sum of the two counts represented the total number of mf ingested by an individual mosquito. To ensure accuracy, counts were rechecked by more than one observer.

Statistical Analyses. Microfilarial counts for both numbers ingested and numbers penetrating the midgut were not normally distributed (Shapiro–Wilk normality tests; W statistic ≥ 0.63 ; $P < 0.0001$) nor did transformation of data by logarithms or square roots normalize the data. Kruskal–Wallis tests were used to compare medians. Infection prevalences were compared using chi-square or Fisher exact tests. Spearman rank correlation (r_s) was used to test for density relationships. A 0.05 level of significance was used for all statistical tests (Statistix version 8, Analytical Software, Tallahassee, FL).

Results

Ingestion of Microfilariae. Based on the results from two volunteers that were fed upon twice, there were no differences among mosquitoes between nighttime and daytime feedings with respect to either quantities of mf ingested (Wilcoxon rank sum tests, $P > 0.27$) or prevalences of mf ingested (Fisher exact tests, $P > 0.63$) (data not shown). Because there was no evidence to suggest that *M. ozzardi* in Trinidad were nocturnally periodic, data for the different times of day were pooled for each of these volunteers in subsequent analyses. Microfilaremiias in the six volunteers ranged from 3 to 394 mf per 200 μ l of blood (Table 1). Only three of a total of 87 mosquitoes dissected were

lost to “dirty” dissections. Overall, mf were detected in 63 of the 84 engorged mosquitoes successfully dissected and examined. The percentages of engorged mosquitoes that ingested mf ranged from 0 (volunteer 5) to 94% (volunteer 2) and were positively correlated with microfilaremia of the volunteers ($r_s = 0.83$, $n = 6$, $P = 0.03$). The mean numbers ingested were mostly within the range expected, based on host microfilaremia and bloodmeal volumes, and there was no indication that *Ae. aegypti* concentrated *M. ozzardi* mf during feeding. There was a significant difference among the six volunteers in the quantity of mf ingested by the mosquitoes that fed on them (range, 0–8.6 mf per mosquito, Kruskal–Wallis, $H = 27.1$, $P < 0.0001$) and in general, the quantity ingested increased with increasing host microfilaremia, although the correlation was not statistically significant ($r_s = 0.81$, $n = 6$, $P = 0.06$).

Mf Penetration of the Mosquito Midgut. Only mosquitoes that had ingested mf were included in the analyses of mf penetration of the mosquito midgut. Overall, *M. ozzardi* mf penetrated the midguts in 43% of 63 mosquitoes that ingested mf (Table 2). Of the 539 total mf ingested by the 63 mosquitoes, 11% penetrated the midgut and moved into the hemocoel. There was a significant positive correlation between the numbers of mf ingested and the numbers of mf that penetrated the midgut ($r_s = 0.45$, $n = 63$, $P < 0.001$) (Fig. 1). Most mf found within mosquito hemocoels were located in the vicinity of the thoracic flight muscles. Examination of mf penetration at selected intervals indicated that mf penetration of the mosquito midgut increased over the first 2 h, remained relatively constant up to 6 h, and then decreased at 14–17 h (Table 3). Unlike mf observed wriggling in the hemo-

Table 2. Penetration of ingested *M. ozzardi* mf through the midgut of resident *Ae. aegypti* mosquitoes fed on microfilaremic human volunteers, Blanchisseuse Trinidad, 2004

Volunteer no.	No. mosquitoes ingesting mf	No. mosquitoes with mf in the hemocoel (%)	Total no. mf ingested	Total no. mf in hemocoel	Penetration efficiency
2	33	16 (48)	453	48	11
3	19	10 (53)	64	13	20
4	6	0 (0)	17	0	0
6	2	0 (0)	2	0	0
1	3	1 (33)	3	1	33
Total	63	27 (43)	539	62	11

Only mosquitoes that ingested mf are included.




Fig. 1. Density relationship between the number of *M. ozzardi* mf ingested by *Ae. aegypti* mosquitoes and the number of mf that penetrated the mosquito midgut and move into the hemocoel ($n = 63$). Some mosquitoes yielded the same values for mf ingested and mf escaped. Thus, some data points have associated superscripts that indicate the number of mosquitoes having those values in common.

coel during earlier time points, most mf observed in the hemocoel after 14 h were immobile and assumed a rigidly straight body position. Melanized mf were never observed in either the midguts or the hemocoel.

Discussion

Mosquitoes are not considered to be involved in the transmission of *M. ozzardi*. Nevertheless, we demonstrated that ingested *M. ozzardi* mf readily penetrated the midguts of native *Ae. aegypti* mosquitoes. Virtually all mf that had penetrated the midguts were either moribund or dead by 14 h after the bloodmeal, confirming that *Ae. aegypti* are not likely to be involved in the natural transmission of this filarial worm. Thus, the importance of this study has nothing to do with transmission of *M. ozzardi*, but rather it addresses the broader question of whether a nonmosquito borne filarial species, like *M. ozzardi*, could enhance the transmission of arboviruses from a vertebrate host (e.g., human) to mosquitoes (e.g., *Ae. aegypti*).

It has been shown in laboratory studies that mosquitoes that feed on vertebrates co-infected with mf and an arbovirus are significantly more efficient vectors of the virus than mosquitoes that ingest virus alone (Turell et al. 1984, 1987; Zytoon et al. 1993; Vaughan and Turell 1996, Vaughan et al. 1999). For this en-

hancing effect to happen, the mf must be able to penetrate the mosquito's midgut. To our knowledge, the only study examining *M. ozzardi* in mosquitoes was by Davis (1928) who dissected triatomine bugs and mosquitoes at various times after feeding them on microfilaremic people and examined the insects for developing filariae in the hemocoel. A small percentage of some mosquito species (including *Ae. aegypti*), contained stunted, developmentally arrested, or dead worms in the thorax. Once it was discovered that *Culicoides* were the vectors (Buckley 1934), no further studies were published on possible relationships between *M. ozzardi* and mosquitoes. Our study corroborates and expands that of Davis (1928) and shows that *M. ozzardi* mf are ingested during the course of blood feeding by *Ae. aegypti* mosquitoes and are capable of penetrating the midgut and moving into the hemocoel. Thus, the potential exists for *M. ozzardi* to be involved in the mf enhancement of anthroponotic viruses such as the dengue viruses.

There are other reasons to suspect that *M. ozzardi* infections could potentially interact with dengue transmission. Throughout most of its distribution in the Western Hemisphere, dengue is sympatric with *M. ozzardi*. Therefore, it is not unreasonable to assume that dually infected humans may exist wherever dengue and mansonelliasis overlap, particularly if the prevalence of *M. ozzardi* infection is high. In Trinidad, prevalence of mansonelliasis has been recorded as high as 11–31% (Nathan et al. 1979, Gonzalez et al. 1999). Mansonelliasis is typically a rural disease due to the distribution of its vector (i.e., biting midges), whereas dengue tends to be a more urban disease because its principal vector, *Ae. aegypti*, is synanthropic. However, in Trinidad, *Ae. aegypti* abundance (as measured by Breteau index) and dengue incidence (i.e., per capita number of new cases) have been observed to be distributed equally between rural and urban areas during recent dengue outbreaks (Brown et al. 2004, Chadee et al. 2005). In addition, the mf of *M. ozzardi* are aperiodic (Nathan et al. 1978). Thus, *M. ozzardi* mf are present in human peripheral circulation during the time when *Ae. aegypti* mosquitoes are most likely to feed (i.e., during daytime), unlike the mf of most strains of *Wuchereria bancrofti* and *Brugia* spp. that display nocturnal periodicity.

It is still unknown whether *M. ozzardi* infections can actually enhance dengue transmission. Proof will require vector competence studies by using *M. ozzardi* mf, similar to studies described by Turell et al. 1984, who demonstrated enhancement of dengue virus infectivity to *Ae. aegypti* by *B. malayi* mf. We hypothesize that the primary effect of mf enhancement on dengue transmission will be a reduction in the extrinsic incubation period of dengue viruses within mosquito species that are already vectors of dengue (e.g., *Ae. aegypti* and *Aedes albopictus* Skuse). All evidence to date indicates that intrathoracically inoculated dengue viruses only replicate efficiently in *Aedes* spp. mosquitoes (Rosen et al. 1985, Huang et al. 1992). Therefore, it seems unlikely that in the case of dengue, mf enhancement can alter the vector competence of

Table 3. Penetration of ingested *M. ozzardi* mf through the midgut of resident *Ae. aegypti* mosquitoes at selected intervals after feeding on microfilaremic human volunteers, Blanchisseuse Trinidad, 2004.

Time after feeding (h)	% of mosquitoes whose midguts were penetrated by ingested mf (no. mosquitoes ingesting mf)	% of ingested mf that penetrated the midgut (no. of mf ingested)
1	14 (7)	5 (20)
2–3	54 (11)	14 (184)
4–6	63 (19)	16 (122)
14–17	32 (25)	7 (213)

Only mosquitoes that ingested mf are included.

otherwise refractory mosquito species, i.e., mf enhancement probably will not create new dengue vectors. Even so, shortening the extrinsic incubation period of dengue within the vector may still produce a significant impact on dengue epidemiology.

Acknowledgments

We thank Robin Persad and Lester James, Ministry of Health, Saint Joseph, Trinidad and Tobago, for collecting and rearing of mosquitoes and invaluable help with the logistics of this study. Nathan Mickelson, University of North Dakota, assayed mosquitoes for flaviviruses. This project was funded in part by a University of North Dakota Faculty Research Seed Money Award (to J.A.V.) and North Dakota EPSCoR (to J.A.B.).

References Cited

- Bell, J. A., N. J. Mickelson, and J. A. Vaughan. 2005. West Nile virus in host-seeking mosquitoes from a residential neighborhood in Grand Forks, North Dakota. *Vector Borne Zoonotic Dis.* 5: 373–382.
- Brown, T. U., K. Babb, M. Nimrod, C.V.F. Carrington, R. A. Salas, and M. A. Monteil. 2004. A retrospective study of the 1996 DEN-1 epidemic in Trinidad: demographic and clinical features. *Dengue Bull.* 28: 7–19.
- Buckley, J.J.C. 1934. On the development, in *Culicoides furcens* Poey, of *Filaria* (= *Mansonella*) *ozzardi* Manson, 1897. *J. Helminthol.* 12: 99–118.
- Calheiros, M. L., G. Fontes, P. Williams, and E.M.M. Rocha. 1998. Experimental infection of *Culex* (*Culex*) *quinquefasciatus* and *Aedes* (*Stegomyia*) *aegypti* with *Wuchereria bancrofti*. *Mem. Inst. Oswaldo Cruz* 93: 855–860.
- Chadee, D. D., C. C. Tilluckdharry, S. C. Rawlins, R. Doon, and M. B. Nathan. 1995. Mass chemotherapy with diethylcarbamazine for the control of Bancroftian filariasis: twelve year follow-up in northern Trinidad, including observations on *Mansonella ozzardi*. *Am. J. Trop. Med. Hyg.* 52: 174–176.
- Chadee, D. D., F.L.R. Williams, and U. D. Kitron. 2005. Impact of vector control on a dengue fever outbreak in Trinidad, West Indies, in 1998. *Trop. Med. Intl. Health* 10: 748–754.
- Chamberlain, R. W., and W. D. Sudia. 1961. Mechanism of transmission of viruses by mosquitoes. *Annu. Rev. Entomol.* 6: 371–390.
- Clements, A. N. 1992. The biology of mosquitoes. vol. 1. Development, nutrition and reproduction. Chapman & Hall, New York.
- Davis, N. C. 1928. A study on the transmission of filaria in northern Argentina. *Am. J. Hyg.* 8: 457–466.
- Gonzalez, A. A., D. D. Chadee, and S. C. Rawlins. 1999. Ivermectin treatment of mansonellosis in Trinidad. *West Indian Med. J.* 48: 231–234.
- Huang, G., E. Vergne, and D. J. Gubler. 1992. Failure of dengue viruses to replicate in *Culex quinquefasciatus* (Diptera: Culicidae). *J. Med. Entomol.* 29: 911–914.
- Kramer, L. D., J. L. Hardy, S. B. Presser, and E. J. Houk. 1981. Dissemination barriers for western equine encephalomyelitis virus in *Culex tarsalis* infected after ingestion of low viral doses. *Am. J. Trop. Med. Hyg.* 30: 190–197.
- Lowichik, A., and R. C. Lowrie. 1988. Uptake and development of *Wuchereria bancrofti* in *Aedes aegypti* and Haitian *Culex quinquefasciatus* that were fed on a monkey with low-density microfilaremia. *Trop. Med. Parasit.* 39: 227–229.
- MacDonald, G. 1952. The analysis of the sporozoite rate. *Trop. Dis. Bull.* 49: 569–586.
- McGreevy, P. B., N. Kostrup, J. Tao, M. M. McGreevy, and T. F. de Marshall. 1982. Ingestion and development of *Wuchereria bancrofti* in *Culex quinquefasciatus*, *Anopheles gambiae* and *Aedes aegypti* after feeding on humans with varying densities of microfilariiae in Tanzania. *Trans. R. Soc. Trop. Med. Hyg.* 76: 288–296.
- Mellor, P. S., and J. Boorman. 1980. Multiplication of blue-tongue virus in *Culicoides nubeculosus* (Meigen) simultaneously infected with the virus and the microfilariiae of *Onchocerca cervicalis* (Railliet & Henry). *Ann. Trop. Med. Parasitol.* 74: 463–469.
- Nathan, M. B. 1978. *Culicoides phlebotomus*, a vector of *Mansonella ozzardi* in coastal north Trinidad, West Indies. *Trans. R. Soc. Trop. Med. Hyg.* 72: 436–437.
- Nathan M. B., C. F. Bartholomew, and E. S. Tikasingh. 1978. The detection of *Mansonella ozzardi* microfilariiae in the skin and blood with a note on the absence of periodicity. *Trans. R. Soc. Trop. Med. Hyg.* 72: 420–422.
- Nathan, M. B., E. S. Tikasingh, G. S. Nelson, A. Santiago, and J. B. Davies. 1979. The prevalence and distribution of *Mansonella ozzardi* in coastal north Trinidad, W.I. *Trans. R. Soc. Trop. Med. Hyg.* 73: 299–302.
- Nathan, M. B., G. Beckles, E. S. Tikasingh, P. J. Hamilton, and S. Monteil. 1982. Parasitological and clinical studies of *Wuchereria bancrofti* and *Mansonella ozzardi* in coastal north Trinidad, West Indies. *West Indies Med. J.* 31: 168–176.
- Nelson, G. S., and J. B. Davies. 1976. Observations on *Mansonella ozzardi* in Trinidad. *Trans. R. Soc. Trop. Med. Hyg.* 70: 16–17.
- Ramachandran, C. P., and M. A. Zaini. 1967. Studies on the transmission of sub-periodic *Brugia malayi* by *Aedes* (*Finlaya*) *togoi* in the laboratory. I. The intake and migration of microfilariiae. *Med. J. Malaya* 22: 136–144.
- Rosen, L., L. E. Roseboom, D. J. Gubler, J. C. Lien, and B. N. Chanotis. 1985. Comparative susceptibility of mosquito species and strains to oral and parenteral infection with dengue and Japanese encephalitis viruses. *Am. J. Trop. Med. Hyg.* 34: 603–615.
- Scaramozzino, N., J. Crance, A. Jouan, D. A. DeBriel, F. Stoll, and D. Garin. 2001. Comparison of Flavivirus universal primer pairs and development of a rapid, highly sensitive heminested reverse transcriptase-PCR assay for detection of flaviviruses targeted to a conservative region of the NS5 gene sequences. *J. Clin. Microbiol.* 39: 1922–1927.
- Shelley, A. J., A.P.A. Luna Dias, and M.A.P. Moraes. 1980. *Simulium* species of the *amazonicum* group as vectors of *Mansonella ozzardi* in the Brazilian Amazon. *Trans. Royal Soc. Trop. Med. Hyg.* 74: 794–788.
- Shelley, A. J., and S. Coscaron. 2001. Simuliid blackflies (Diptera: Simuliidae) and ceratopogonid midges (Diptera: Ceratopogonidae) as vectors of *Mansonella ozzardi* (Nematoda: Onchocercidae) in northern Argentina. *Mem. Inst. Oswaldo Cruz* 96: 451–458.
- Turell, M. J., P. A. Rossignol, A. Spielman, C. A. Rossi, and C. L. Bailey. 1984. Enhanced arboviral transmission by mosquitoes that concurrently ingested microfilariiae. *Science* (Wash., D.C.) 225: 1039–1041.
- Turell, M. J., T. N. Mather, A. Spielman, and C. L. Bailey. 1987. Increased dissemination of dengue 2 virus in *Aedes aegypti* associated with concurrent ingestion of microfilariiae of *Brugia malayi*. *Am. J. Trop. Med. Hyg.* 37: 197–201.

- Vaughan, J. A., and M. J. Turell. 1996. Dual host infections: enhanced infectivity of eastern equine encephalitis virus to *Aedes* mosquitoes mediated by *Brugia* microfilariae. *Am. J. Trop. Med. Hyg.* 54: 105–109.
- Vaughan, J. A., M. Trpis, and M. J. Turell. 1999. *Brugia malayi* microfilariae enhance the infectivity of Venezuelan equine encephalitis virus to *Aedes* mosquitoes. *J. Med. Entomol.* 36: 758–763.
- Zielke, E. 1992. Observations on the midgut penetration by *Wuchereria bancrofti* microfilariae in mosquitoes of varying susceptibilities to infection. *Mosquito-borne Dis. Bull.* 9: 60–63.
- Zytoon, E. M., H. I. El-Belbasi, and T. Matsumura. 1993. Mechanism of increased dissemination of chikungunya virus in *Aedes albopictus* mosquitoes concurrently ingesting microfilariae of *Dirofilaria immitis*. *Am. J. Trop. Med. Hyg.* 49: 201–207.

Received 5 July 2006; accepted 12 October 2006.
